


LidoDose™

Now Available

The Single Dose™ Topical Anesthetic & Antiseptic for Preprocedural Use.

For preparation of the skin prior to an injection. For pain associated with any procedure requiring penetration of the skin, such as dialysis, vaccinations, venipuncture and other procedures.



Suitable for Use on Children Ages Two and Above.

Works fast.

Onset of action as fast as 5 minutes from time of application.



Antimicrobial.

Kills >99.99% of common skin pathogens within 3 minutes.

Care Settings.

Acute
Subacute/LTC
Infusion
Ambulatory
Physician Offices
Clinics



READ THIS INFORMATION BEFORE USING THIS PRODUCT

DOSAGE AND ADMINISTRATION:

Adults and children 2 years of age and older: Apply using a gloved hand. Swirl the applicator tip prior to removing swab from the pouch and apply to area 3-5 minutes prior to procedure not more than 4 times daily.

Children under 2 years of age: Consult a physician.

WARNING:

For External Use Only. Avoid contact with eyes.

Stop use and consult a physician if condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days.

Do not use in large quantities, particularly over raw surfaces or blistered areas.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

STORAGE:

Store between 68° and 77°F (20-25°C), with allowed excursions between 59° and 86°F (15-30°C). Protect from freezing [See USP Controlled Room Temperature].

ACTIVE INGREDIENTS:

Lidocaine Hydrochloride 3%.....Anesthetic

INACTIVE INGREDIENTS:

Aqua (Deionized Water), Carbomer, Isopropyl Alcohol, Petrolatum, Polysorbate-20, Triethanolamine

Manufactured for
Gensco Pharma
Miami, FL 33122
855-743-6726

www.genscopharma.com



Drug Facts

Active Ingredient	Purpose
Lidocaine Hydrochloride 3%	Topical anesthetic

Uses Temporary relief of minor pain.

Warnings For external use only.
Avoid contact with eyes.

When using this product

- do not use in large quantities, particularly over raw surfaces or blistered areas

Stop use and ask doctor

- if condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions Apply using a gloved hand. Swirl the applicator tip prior to removing swab from the pouch.

Adults and children over 2 years of age	Apply to area 3-5 minutes prior to procedure not more than 4 times daily.
---	---

Children under 2	Consult a physician
------------------	---------------------

Other information

- package not child resistant
- avoid storing product in direct sunlight
- protect product from excessive moisture
- store between 68° and 77°F (20-25°C)

Inactive ingredients Aqua (Deionized Water), Carbomer, Isopropyl Alcohol, Petrolatum, Polysorbate-20, Triethanolamine

Questions or Comments?

1-855-743-6726.